

In order to receive full credit, SHOW ALL YOUR WORK. Full credit will be given only if all reasoning and work is provided. When applicable, please enclose your final answers in boxes.

1. (10 Points) True/False and Short Response

(a) Mark each statement True or False.

i. The columns of a matrix  $A$  are linearly independent if the equation  $Ax = 0$  has only the trivial solution,  $x = 0$ . True, Col. lin. ind.  $\Rightarrow$  pivot in every column  $\Rightarrow$

$\Rightarrow$  no free vars  $\Rightarrow \vec{x} = \vec{0}$  is the only soln.

ii. If the equation  $Ax = 0$  has a nontrivial solution, then  $A$  has fewer than  $n$  pivot positions.

True. See above

iii. If the columns of  $A$  are linearly dependent, then  $\det(A) = 0$ .

lin dep.  $\Rightarrow$  fewer than  $n$ -pivots  $\Rightarrow A \neq I \Rightarrow A^{-1}$  DNE  $\Rightarrow \det(A) = 0$

iv. The columns of an  $n \times n$  invertible matrix form a basis for  $\mathbb{R}^n$ .

$n \times n$  invert  $\Rightarrow \det(A) \neq 0 \Rightarrow n$ -pivots  $\Rightarrow A \sim I \Rightarrow$  Col. Basis for  $\mathbb{R}^n$   
True

v. The column space of  $A$  is the set of all solutions to  $Ax = b$ .

False. Column Space: Set of all  $\sum x_i \vec{a}_i$   
Set of Soln is Set of all  $\vec{x}$  which solves  $A\vec{x} = \vec{b}$

(b) Please respond to one of the following:

i. Suppose  $A$  is a  $4 \times 3$  matrix and  $b$  is a vector in  $\mathbb{R}^4$  with the property that  $Ax = b$  has a unique solution. What can you say about the reduced echelon form of  $A$ ?

ii. Suppose  $A = \begin{bmatrix} 0 & 0 & 0 \\ 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$  what is the determinant of  $A$ ? List three more equivalent properties/characterizations of  $A$ .

iii. Suppose that  $Ax = 0$  and  $x \neq 0$  is an eigenvector. What does the eigenvalue corresponding to this  $x$  have to be? Explain.

i)  $A \sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$

ii)  $A^{-1}$  does not exist,  $A \neq I$ ,  $A\vec{x} = \vec{0}$  has nonunique soln  
 $\det(A) = 0$

iii)  $\vec{x}$  must satisfy  $A\vec{x} = \lambda\vec{x}$

but  $A\vec{x} = 0 \Rightarrow \lambda\vec{x} = 0 \Rightarrow \lambda = 0$  is its associated Eigenvalue

2. (10 Points) Given the following matrix  $A$  and its associated eigenvectors  $y_1$  and  $y_2$ :

$$A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} \quad y_1 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \quad y_2 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad b_1 = \begin{bmatrix} 0 \\ 0 \end{bmatrix}, \quad b_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad b_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}.$$

(a) Find the solution to  $Ax = b_1$ .

$$\left[ \begin{array}{cc|c} 1 & -1 & 0 \\ -1 & 1 & 0 \end{array} \right] \sim \left[ \begin{array}{cc|c} 1 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow x_1 = x_2 \Rightarrow \bar{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} x_2, \quad x_2 \in \mathbb{R}$$

(b) Find the solution to  $Ax = b_2$ .

$$\left[ \begin{array}{cc|c} 1 & -1 & 1 \\ -1 & 1 & -1 \end{array} \right] \sim \left[ \begin{array}{cc|c} 1 & -1 & 1 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow \bar{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} x_2 + \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad x_2 \in \mathbb{R}$$

(c) Find the solution to  $Ax = b_3$ .

$$\left[ \begin{array}{cc|c} 1 & -1 & 2 \\ -1 & 1 & 2 \end{array} \right] \sim \left[ \begin{array}{cc|c} 1 & -1 & 1 \\ 0 & 0 & 4 \end{array} \right] \Rightarrow \text{No Soln}$$

(d) Find  $A^4$

$$\begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}^4 \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix} = \begin{bmatrix} -2^4 & 0 \\ 2^4 & 0 \end{bmatrix} \begin{bmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix} = \begin{bmatrix} \frac{2^4}{2} & -\frac{2^4}{2} \\ -\frac{2^4}{2} & \frac{2^4}{2} \end{bmatrix} = \begin{bmatrix} 2^3 & -2^3 \\ -2^3 & 2^3 \end{bmatrix}$$

$2^3 = 8$

3. (10 Points) Given,

$$\begin{aligned} x_1 + hx_2 &= 2 \\ 4x_1 + 8x_2 &= k. \end{aligned} \quad \left[ \begin{array}{cc|c} 1 & h & 2 \\ 4 & 8 & k \end{array} \right] = \left[ \begin{array}{cc|c} 1 & h & 2 \\ 0 & 8-4h & k-8 \end{array} \right]$$

Choose  $h$  and  $k$  such that the system has:

(a) No Solution

$$h=2, \quad k \neq 8$$

(b) A Unique Solution

$$h \neq 2, \quad k \in \mathbb{R}$$

(c) Many Solutions

$$h=2, \quad k=8$$

4. (10 Points) Given,

$$A = \begin{bmatrix} 1 & 2 & -5 & 11 & -3 \\ 2 & 4 & -5 & 15 & 2 \\ 1 & 2 & 0 & 4 & 5 \\ 3 & 6 & -5 & 19 & -2 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 0 & 4 & 5 \\ 0 & 0 & 5 & -7 & 8 \\ 0 & 0 & 0 & 0 & -9 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Find a basis and the dimension of the null-space and column-space of A.

$$\vec{x} = \begin{bmatrix} -2x_2 - 4x_4 \\ x_2 \\ \frac{7}{5}x_4 \\ x_4 \\ 0 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} -4 \\ 0 \\ \frac{7}{5} \\ 1 \\ 0 \end{bmatrix}, x_2, x_4 \in \mathbb{R}$$

$$B_{\text{Null}(A)} = \left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ 0 \\ \frac{7}{5} \\ 1 \\ 0 \end{bmatrix} \right\}$$

$$\dim(B_{\text{Null}(A)}) = 2$$

$$B_{\text{Col}(A)} = \left\{ \begin{bmatrix} 1 \\ 2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} -5 \\ -5 \\ 0 \\ -5 \end{bmatrix}, \begin{bmatrix} -3 \\ 2 \\ -5 \\ -2 \end{bmatrix} \right\} \quad \dim[B_{\text{Col}(A)}] = 3$$

5. (10 Points) Given,

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

Find all eigenvalues and eigenvectors of A.

$$\lambda = 1, 1, -1$$

$$\lambda = 1$$

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -2 \end{bmatrix} \Rightarrow \vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ 0 \end{bmatrix} \Rightarrow \vec{x}^{(1)} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \vec{x}^{(2)} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$\lambda = -1$$

$$\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix} \Rightarrow \vec{x}^{(3)} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$