

In order to receive full credit, **SHOW ALL YOUR WORK**. Full credit will be given only if all reasoning and work is provided. When applicable, please enclose your final answers in boxes.

1. (10 Points)

(a) True/False: No Justification Needed

- i. The function e^{ix} has no symmetry.
- ii. If a periodic function is neither even nor odd then its Fourier series representation must have sine terms/modes.
- iii. If the complex Fourier coefficients are purely imaginary then the periodic function is even.
- iv. A function can have only one periodic extension.
- v. A truncated Fourier sine half-range expansion will not have Gibb's phenomenon.

(b) Short Response

- i. Provide two physical interpretations of both Fourier coefficients and their corresponding Fourier modes.
- ii. Explain Gibb's phenomenon. What is it and when can you expect it to occur?

2. (10 Points)

(a) Given that n is an integer for following integrals,

$$\int_{-\pi}^{\pi} f(x) dx = \frac{x^3}{3} \Big|_{-\pi}^{\pi},$$

$$\int_{-\pi}^{\pi} f(x) \cos(nx) dx = \left[\frac{x^2 \sin(nx)}{n} + \frac{2x \cos(nx)}{n^2} - \frac{2 \sin(nx)}{n^3} \right] \Big|_{-\pi}^{\pi},$$

$$\int_{-\pi}^{\pi} f(x) \sin(nx) dx = \left[\frac{x^2 \cos(nx)}{n} + \frac{2x \sin(nx)}{n^2} - \frac{2 \cos(nx)}{n^3} \right] \Big|_{-\pi}^{\pi},$$

$$\int_{-\pi}^{\pi} g(x) dx = e^{-i\pi} - e^{i\pi},$$

$$\int_{-\pi}^{\pi} g(x) e^{-inx} dx = e^{inx} \left(\frac{1}{n^2} - \frac{ix}{n} \right) \Big|_{-\pi}^{\pi}.$$

i. Calculate the symmetry and Fourier series of $f(x)$.


ii. Calculate the symmetry and Fourier series of $g(x)$.

iii. From the complex Fourier series of g calculate the corresponding real Fourier series.

(b) The following table contains different boundary conditions for the ODE, $F'' + \lambda F = 0$, $\lambda \in [0, \infty)$. Fill in each table element with either a yes or a no.

	Boundary value problem has a cosine solution	Boundary value problem has a sine solution	Boundary value problem has a nontrivial constant solution
$F(0) = 0, F'(L) = 0$			
$F'(0) = 0, F'(L) = 0$			
$F(0) = 0, F(L) = 0$			
$F'(0) = 0, F(L) = 0$			

3. (10 Points) Suppose f is given by the graph below.



- (a) On the graph above, sketch the Fourier cosine and sine half-range expansions with dashed lines and solid lines, respectively.
- (b) Find the Fourier coefficients of the cosine and sine half-range expansions. Justify your calculations.

4. (10 Points) Find the complex Fourier series representation of

$$f(x) = \begin{cases} 0, & -1 < x < 0, \\ x, & 0 < x < 1. \end{cases}$$

5. (10 Points) Given,

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}, \quad x \in (0, \pi), \quad t \in (0, \infty), \quad (1)$$

$$u_x(0, t) = 0, \quad u_x(\pi, t) = 0, \quad (2)$$

$$u(x, 0) = f(x). \quad (3)$$

(a) Describe the separation of variable procedure used to solve the partial differential equation given by (1)-(3). Be sure to discuss how each step corresponds to each equation (1), (2) and (3).

(b) The separation of variables process, applied to (1), yields the equation

$$F''(x) + \lambda F(x) = 0, \quad \lambda \in \mathbb{R}. \quad (4)$$

Find all nontrivial solutions of (4) that satisfies (2). Justify your choices.