

Quote of Homework: Linear Algebra Part I

**Paul Atreides:** Fear is the mind-killer. Fear is the little-death that brings total obliteration.

Frank Herbert : Dune (1965)

## 1. MATRIX MULTIPLICATION

Define the *commutator* and *anti-commutator* of two square matrices to be,

$$[\cdot, \cdot] : \mathbb{C}^{n \times n} \times \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}, \text{ such that } [\mathbf{A}, \mathbf{B}] = \mathbf{AB} - \mathbf{BA}, \text{ for all } \mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n},$$

$$\{\cdot, \cdot\} : \mathbb{C}^{n \times n} \times \mathbb{C}^{n \times n} \rightarrow \mathbb{C}^{n \times n}, \text{ such that } \{\mathbf{A}, \mathbf{B}\} = \mathbf{AB} + \mathbf{BA}, \text{ for all } \mathbf{A}, \mathbf{B} \in \mathbb{C}^{n \times n},$$

respectively. Also define the *Kronecker delta* and *Levi-Civita* symbols to be,

$$\delta_{ij} : \mathbb{N} \times \mathbb{N} \rightarrow \{0, 1\}, \text{ such that } \delta_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j \end{cases}$$

$$\epsilon_{ijk} : (i, j, k) \rightarrow \{-1, 0, 1\}, \text{ such that } \epsilon_{ijk} = \begin{cases} 1, & \text{if } (i, j, k) \text{ is } (1, 2, 3), (2, 3, 1) \text{ or } (3, 1, 2), \\ -1, & \text{if } (i, j, k) \text{ is } (3, 2, 1), (1, 3, 2) \text{ or } (2, 1, 3), \\ 0, & \text{if } i = j \text{ or } j = k \text{ or } k = i \end{cases}$$

respectively. Also define the so-called *Pauli spin-matrices* (PSM) to be,

$$\sigma_1 = \sigma_x = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \sigma_2 = \sigma_y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}, \quad \sigma_3 = \sigma_z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}.$$

1.1. **The PSM are *self-adjoint* matrices.** Show that  $\sigma_m = \sigma_m^H$  for  $m = 1, 2, 3$ .

1.2. **The PSM are *unitary* matrices.** Show that  $\sigma_m^2 = \mathbf{I}$  for  $m = 1, 2, 3$  where  $[\mathbf{I}]_{ij} = \delta_{ij}$ .

1.3. **Trace and Determinant.** Show that  $\text{tr}(\sigma_m) = 0$  and  $\det(\sigma_m) = -1$  for  $m = 1, 2, 3$ .

1.4. **Anti-Commutation Relations.** Show that  $\{\sigma_i, \sigma_j\} = 2\delta_{ij}\mathbf{I}$  for  $i = 1, 2, 3$  and  $j = 1, 2, 3$ .

1.5. **Commutation Relations.** Show that  $[\sigma_i, \sigma_j] = 2\sqrt{-1} \sum_{k=1}^3 \epsilon_{ijk} \sigma_k$  for  $i = 1, 2, 3$  and  $j = 1, 2, 3$ .

## 2. SOLUTIONS SETS TO LINEAR SYSTEMS OF ALGEBRAIC EQUATIONS

Given,

$$\mathbf{A}_1 = \begin{bmatrix} 1 & -3 & 0 \\ -1 & 1 & 5 \\ 0 & 1 & 1 \end{bmatrix}, \quad \mathbf{A}_2 = \begin{bmatrix} 6 & 18 & -4 \\ -1 & -3 & 8 \\ 5 & 15 & -9 \end{bmatrix}, \quad \mathbf{A}_3 = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 1 & 0 & 3 \end{bmatrix}, \quad \mathbf{A}_4 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 9 \end{bmatrix}, \quad \mathbf{A}_5 = \begin{bmatrix} 5 & 3 \\ -4 & 7 \\ 9 & -2 \end{bmatrix},$$

$$\mathbf{b}_1 = \begin{bmatrix} 5 \\ 2 \\ 0 \end{bmatrix}, \quad \mathbf{b}_2 = \begin{bmatrix} 20 \\ 4 \\ 11 \end{bmatrix}, \quad \mathbf{b}_3 = \begin{bmatrix} 4 \\ 1 \\ 0 \end{bmatrix}, \quad \mathbf{b}_4 = \begin{bmatrix} 10 \\ 20 \\ 30 \end{bmatrix}, \quad \mathbf{b}_5 = \begin{bmatrix} 22 \\ 20 \\ 15 \end{bmatrix}.$$

2.1. **Algebra.** Find all solutions to  $\mathbf{A}_i \mathbf{x} = \mathbf{b}_i$  for  $i = 1, 2, 3, 4, 5$ .

2.2. **Geometry.** Describe or plot the geometry formed by the linear systems and their solution sets.

## 3. SQUARE COEFFICIENT DATA AND MATRIX INVERSION

Given,

$$\mathbf{A} = \begin{bmatrix} 3 & 6 & 7 \\ 0 & 2 & 1 \\ 2 & 3 & 4 \end{bmatrix}.$$

3.1. **Matrix Inverse: Take One.** Find  $\mathbf{A}^{-1}$  using the Gauss-Jordan Method. (pg.317)

3.2. **Matrix Inverse: Take Two.** Find  $\mathbf{A}^{-1}$  using the cofactor representation. (Theorem 2 pg.318)

3.3. **Solutions to Linear Systems.** Using  $\mathbf{A}^{-1}$  find the unique solution to  $\mathbf{Ax} = \mathbf{b} = [b_1 \ b_2 \ b_3]^T$ .

## 4. DETERMINANTS

Given,

$$\mathbf{A} = \begin{bmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{bmatrix}.$$

4.1. **Vandermonde Determinant.** Show that the  $\det(\mathbf{A}) = (c-a)(c-b)(b-a)$ .

4.2. **Application.** Determine which of the following sets of points can be uniquely interpolated by the polynomial  $p(t) = a_0 + a_1t + a_2t^2$ .

$$S_1 = \{(1, 12), (2, 15), (3, 16)\}$$

$$S_2 = \{(1, 12), (1, 15), (3, 16)\}$$

$$S_3 = \{(1, 12), (2, 15), (2, 15)\}$$

5. ROTATION TRANSFORMATIONS IN  $\mathbb{R}^2$  AND  $\mathbb{R}^3$ 

Given,

$$\mathbf{A}(\theta) = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}.$$

5.1. **The Unit Circle.** Show that the transformation  $\mathbf{A}\hat{\mathbf{i}}$  rotates  $\hat{\mathbf{i}} = [1 \ 0]^T$  counter-clockwise by an angle  $\theta$  and defines a parametrization of the *unit circle*. What matrix would undo this transformation?

5.2. **Determinant.** Show that  $\det(\mathbf{A}) = 1$ .

5.3. **Orthogonality.** Show that  $\mathbf{A}^T \mathbf{A} = \mathbf{A} \mathbf{A}^T = \mathbf{I}$ .

5.4. **Rotations in  $\mathbb{R}^3$ .** Given,

$$\mathbf{R}_1(\theta) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{bmatrix}, \quad \mathbf{R}_2(\theta) = \begin{bmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{bmatrix} \quad \mathbf{R}_3(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Describe the transformations defined by each of these matrices on vectors in  $\mathbb{R}^3$ .