

Jan 18 lec covers section 1-4 and 2-4 along with material on conservation laws which is not in the book (references are given below).



Review for the Monday 1/2 exam (30 minutes):

- one problem on deriving areas or volumes of circles, cylinders, or spheres
- two problems on setting up the integral for the electric field from either a line or surface charge distribution.

You will only have to justify the set-up of the integral with limits correctly in these problems.

### InkSurvey muddy points:

- flux thru straw surfaces:  $\int \vec{E} \cdot d\vec{a} \propto \int \frac{1}{r^2} da \cos \theta$



What happens with a charge offset from the center of the first sphere?

Solid angle: [http://en.wikipedia.org/wiki/Solid\\_angle](http://en.wikipedia.org/wiki/Solid_angle)  $\int d\Omega = \int \sin \theta \, d\theta \, d\phi = 4\pi$

$$-\oint \vec{E} \cdot d\vec{a} = \frac{Q_{\text{enc}}}{\epsilon_0}$$

neg flux      neg charge



How do you get flux from differential form?


$$\vec{\nabla} \cdot \vec{E} = \rho / \epsilon_0$$

$$\int \vec{\nabla} \cdot \vec{E} d\tau = \frac{1}{\epsilon_0} \int \rho d\tau$$

↑ volume dx dy dz

div. theorem

$$\oint \vec{E} \cdot d\vec{a} = \frac{Q_{\text{enc}}}{\epsilon_0}$$



$$\int_a^b \frac{df}{dx} dx = f(b) - f(a)$$

Fundamental theorem of calculus

sum changes in f to get total change at bndry

Integral of derivative over interval is the value of function at end points (bndries)

Fundamental theorem of divergences

$$\int \vec{\nabla} \cdot \vec{F} d\tau = \oint \vec{F} \cdot d\vec{a}$$

Integral of derivative (divergence) over region (volume) is value of function at bndry (surface)

- Analogy questions like

$$\vec{\nabla} \cdot \vec{B} = ? \quad \vec{\nabla} \times \vec{E} = ?$$

$$\oint \vec{B} \cdot d\vec{a} =$$


-Advantages of differential form: Coulombs law becomes a partial differential equation.

$$\vec{\nabla} \cdot \vec{E} = \rho / \epsilon_0$$

Use numerical method to iterate solns

## Examples of Gauss's law

### uniformly charged sphere




$$\oint \vec{E} \cdot d\vec{a} = E 4\pi R^2$$

$$\frac{Q_{\text{enc}}}{\epsilon_0} = \frac{\sigma 4\pi r^2}{\epsilon_0}$$

Close to the surface

### Infinite plane of charge




### Superposition principle

Conservation of charge continued.

$$\oint \vec{J} \cdot d\vec{a} = - \frac{dQ_{\text{enclosed}}}{dt}$$

by this surface



$$Q_{\text{enclosed}} = \int \rho \, d\text{volume}$$

"  $dx dy dz$  or  $r^2 \sin \theta d\theta d\phi dr$

$$- \frac{dQ_{\text{enclosed}}}{dt} = \int - \frac{\partial \rho}{\partial t} dx dy dz$$

$$\oint \vec{J} \cdot d\vec{a} = \int \vec{\nabla} \cdot \vec{J} dx dy dz \quad \text{Divergence theorem}$$

Differential form of conservation of charge is

$$\vec{\nabla} \cdot \vec{J} = - \frac{\partial \rho}{\partial t}$$

Also called the CONTINUITY EQUATION

[http://en.wikipedia.org/wiki/Continuity\\_equation](http://en.wikipedia.org/wiki/Continuity_equation)

this link gives examples for 4-current in special relativity, fluid dynamics, thermodynamics, quantum mechanics, general relativity, quantum chromodynamics

traffic flow continuity equation

Examples of vector calculus in fluids

[http://en.wikipedia.org/wiki/Navier%E2%80%93Stokes\\_equations](http://en.wikipedia.org/wiki/Navier%E2%80%93Stokes_equations)