

Eigenproblems : Eigenvalues, Eigenvectors, Diagonalization, Self-Adjoint Operators

Text: 8.1-8.4

Lecture Slides: 7-8

Quote of Homework Three

Raoul Duke: Nonsense. We came here to find the American Dream, and now we're right in the vortex you want to quit? You must realize that we've found the Main Nerve.

Grisoni and Gilliam : Fear and Loathing in Las Vegas (1998)

1. EIGENVALUES AND EIGENVECTORS

$$\mathbf{A}_1 = \begin{bmatrix} 4 & 0 & 1 \\ -2 & 1 & 0 \\ -2 & 0 & 1 \end{bmatrix}, \quad \mathbf{A}_2 = \begin{bmatrix} 3 & 1 \\ -2 & 1 \end{bmatrix}, \quad \mathbf{A}_3 = \begin{bmatrix} 4 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{bmatrix}, \quad \mathbf{A}_4 = \begin{bmatrix} .1 & .6 \\ .9 & .4 \end{bmatrix}, \quad \mathbf{A}_5 = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix},$$

1.1. **Eigenproblems.** Find all eigenvalues and eigenvectors of \mathbf{A}_i for $i = 1, 2, 3, 4, 5$.

2. DIAGONALIZATION

2.1. **Eigenbasis and Decoupled Linear Systems.** Find the diagonal matrix \mathbf{D}_i and vector $\tilde{\mathbf{Y}}$ that completely decouples the system of linear differential equations $\frac{d\tilde{\mathbf{Y}}_i}{dt} = \mathbf{A}_i \tilde{\mathbf{Y}}_i$ for $i = 3, 4, 5$.

3. REGULAR STOCHASTIC MATRICES

For the *regular stochastic matrix* \mathbf{A}_4 , define its associated steady-state vector, \mathbf{q} , to be such that $\mathbf{A}_4 \mathbf{q} = \mathbf{q}$.

3.1. **Limits of Time Series.** Show that $\lim_{n \rightarrow \infty} \mathbf{A}_4^n \mathbf{x} = \mathbf{q}$ where $\mathbf{x} \in \mathbb{R}^2$ such that $x_1 + x_2 = 1$.

4. ORTHOGONAL DIAGONALIZATION AND SPECTRAL DECOMPOSITION

Recall that if $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$ then their inner-product is defined to be $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^H \mathbf{y} = \bar{\mathbf{x}}^T \mathbf{y}$.

4.1. **Self-Adjointness.** Show that \mathbf{A}_5 is a self-adjoint matrix.

4.2. **Orthogonal Eigenvectors.** Show that the eigenvectors of \mathbf{A}_5 are orthogonal with respect to the inner-product defined above.

4.3. **Orthonormal Eigenbasis.** Using the previous definition for length of a vector and the eigenvectors of the self-adjoint matrix, construct an *orthonormal basis* for \mathbb{C}^2 .

4.4. **Orthogonal Diagonalization.** Show that $\mathbf{U}^H = \mathbf{U}^{-1}$, where \mathbf{U} is a matrix containing the normalized eigenvectors of \mathbf{A}_4 .

4.5. **Spectral Decomposition.** Show that $\mathbf{A}_4 = \lambda_1 \mathbf{x}_1 \mathbf{x}_1^H + \lambda_2 \mathbf{x}_2 \mathbf{x}_2^H$.

5. INTRODUCTION TO SELF-ADJOINT OPERATORS

Let L be a linear transformation defined by,

$$(1) \quad Lu = \frac{1}{w(x)} \left(-\frac{d}{dx} \left[p(x) \frac{du}{dx} \right] + q(x)u \right), \text{ where } x \in (a, b) \text{ such that,}$$

$$(2) \quad k_1 u(a) + k_2 u'(b) = 0$$

$$(3) \quad l_1 u(b) + l_2 u'(b) = 0.$$

Finding all nontrivial eigenfunctions of (1), which satisfy the boundary conditions (2)-(3) is called a *Sturm-Liouville Problem* (SLP).

5.1. **A Simple SLP.** Let $p(x) = 1$, $q(x) = 0$, $w(x) = 1$, $k_1 = l_1 = 1$, $k_2 = l_2 = 0$ and $a = 0$, $b = \pi$. Show that the eigenvalue/eigenfunction pairs to the SLP are defined by $u_n(x) = \sin(\sqrt{\lambda_n}x)$, $\lambda_n = n^2$, for $n = 1, 2, 3, \dots$

5.2. **Orthogonality of Eigenfunctions.** Using the abstract inner-product defined in homework 2 problem 5.2, $\langle f, g \rangle = \int_{-\pi}^{\pi} f(x)g(x)dx$, show that the previous eigenfunctions form an orthogonal set. That is, show that $\langle y_n, y_m \rangle = \pi \delta_{nm}$ for $n = 1, 2, 3, \dots$, and $m = 1, 2, 3, \dots$