

Quote of Homework Five

And the feeling is that there's something wrong, 'cause I can't find the words, and I can't find the songs.

Radiohead : Stop Whispering (1993)

1. CONSERVATION LAWS IN ONE-DIMENSION

Recall that the conservation law encountered during the derivation of the heat equation was given by,

$$(1) \quad \frac{\partial u}{\partial t} = -\kappa \nabla \cdot \phi = -\kappa \operatorname{div}(\phi),$$

which reduces to

$$(2) \quad \frac{\partial u}{\partial t} = -\kappa \frac{\partial \phi}{\partial x}, \quad \kappa \in \mathbb{R}$$

in one-dimension of space.<sup>1</sup> In general, if the function  $u = u(x, t)$  represents the density of a physical quantity then the function  $\phi = \phi(x, t)$  represents its flux. If we assume the  $\phi$  is proportional to the negative gradient of  $u$  then, from (2), we get the one-dimensional heat/diffusion equation.<sup>2</sup>

1.1. **Transport Equation.** Assume that  $\phi$  is proportional to  $u$  to derive, from (2), the convection/transport equation,  $u_t + cu_x = 0 \quad c \in \mathbb{R}$ .

1.2. **General Solution to the Transport Equation.** Show that  $u(x, t) = f(x - ct)$  is a solution to this PDE.

1.3. **Diffusion-Transport Equation.** If both diffusion and convection are present in the physical system then the flux is given by,  $\phi(x, t) = cu - du_x$ , where  $c, d \in \mathbb{R}^+$ . Derive from, (2), the convection-diffusion equation  $u_t + \alpha u_x - \beta u_{xx} = 0$  for some  $\alpha, \beta \in \mathbb{R}$ .

1.4. **Convection-Diffusion-Decay.** If there is also energy/particle loss proportional to the amount present then we introduce to the convection-diffusion equation the term  $\lambda u$  to get the convection-diffusion-decay equation.<sup>3</sup>

1.5. **General Importance of Heat/Diffusion Problems.** Given that,

$$(3) \quad u_t = Du_{xx} - cu_x - \lambda u.$$

Show that by assuming,  $u(x, t) = w(x, t)e^{\alpha x - \beta t}$ , (3) can be transformed into a heat equation on the new variable  $w$  where  $\alpha = c/(2D)$  and  $\beta = \lambda + c^2/(4D)$ .<sup>4</sup>

2. ONE DIMENSIONAL HEAT EQUATION WITH SOURCE TERM

Given,

$$(4) \quad \frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} + F(x, t),$$

where  $x \in (0, L)$  and  $t \in (0, \infty)$ , subject to

$$(5) \quad u_x(0, t) = 0, \quad u_x(L, t) = 0,$$

and

$$(6) \quad u(x, 0) = g(x).$$

<sup>1</sup>When discussing heat transfer this is known as Fourier's Law of Cooling. In problems of steady-state linear diffusion this would be called Fick's First Law. In discussing electricity  $u$  could be charge density and  $q$  would be its flux.

<sup>2</sup>AKA Fick's Second Law associated with linear non-steady-state diffusion.

<sup>3</sup>The  $u_{xx}$  term models diffusion of energy/particles while  $u_x$  models convection,  $u$  models energy/particle loss/decay. The final term should not be surprising! Wasn't the appropriate model for radioactive/exponential decay  $Y' = -\alpha^2 Y$ ?

<sup>4</sup>This shows that the general PDE (3), which models can be solved using heat equation techniques.

2.1. **Cosine Half-Range Expansion.** Let  $F(x, t) = e^{-t} \sin\left(\frac{2\pi}{L}x\right)$  be the heat generation function. Find the Fourier cosine half-range expansion of  $F$ .

2.2. **General Solution.** Using the previous result, solve for  $G_n(t)$  for  $n = 0, 1, 2, 3, \dots$  assuming that  $u(x, t) = G_0(t) + \sum_{n=1}^{\infty} \cos\left(\frac{n\pi}{L}x\right) G_n(t)$ .

2.3. **Fourier Coefficients.** Assuming that  $g(x) = \begin{cases} \frac{2k}{L}x, & 0 < x < \frac{L}{2}, \\ \frac{2k}{L}(L-x), & \frac{L}{2} < x < L \end{cases}$ , solve for any unknown constants associated with the general solution.

### 3. TIME DEPENDENT BOUNDARY CONDITIONS

It makes sense to consider time-dependent interface conditions. That is, (4) and (6) subject to

$$(7) \quad u(0, t) = g(t), \quad u(L, t) = h(t), \quad t \in (0, \infty)$$

Show that this PDE transforms into:

$$(8) \quad \frac{\partial w}{\partial t} = c^2 \frac{\partial^2 w}{\partial x^2} - S_t(x, t) \quad ,$$

$$(9) \quad x \in (0, L), \quad t \in (0, \infty), \quad c^2 = \frac{\kappa}{\rho\sigma}.$$

with boundary conditions and initial conditions,

$$(10) \quad w(0, t) = w(L, t) = 0,$$

$$(11) \quad w(x, 0) = F(x),$$

where  $F(x) = f(x) - S(x, 0)$  and  $S(x, t) = \frac{h(t) - g(t)}{L}x + g(t)$ .<sup>5</sup>

### 4. HEAT EQUATION ON A BOUNDED DOMAIN OF $\mathbb{R}^{2+1}$

Suppose that heat is allowed to flow in an  $x, y$ -plane, of finite area,  $A = L_x L_y$ , that has been insulated in the  $z$ -direction and its perimeter.

4.1. **Separation of Variables.** Find three ODEs consistent with the heat equation modeling the physical situation described above.

4.2. **Boundary Value Problems.** Write down the boundary conditions implied by the physical situation above and solve all ODEs, with their corresponding boundary conditions, given by the separation of variables above.

4.3. **Fourier Synthesis.** Apply superposition to the solutions of the ODE/BVPs from the previous step to find the general solution to the heat equation. From the general solution, show that the long-time behavior is to average the initial condition over the plane.

### 5. FOURIER TRANSFORMS

5.1. **Dirac-Delta.**  $\mathfrak{F}\{f\}$  where  $f(x) = \delta(x - x_0)$ ,  $x_0 \in \mathbb{R}$ .<sup>6</sup>

5.2. **Decaying Exponential Function.**  $\mathfrak{F}\{f\}$  where  $f(x) = e^{-k_0|x|}$ ,  $k_0 \in \mathbb{R}^+$ .

5.3. **Even Combination of Dirac-Deltas.**  $\mathfrak{F}^{-1}\{\hat{f}\}$  where  $\hat{f}(\omega) = \frac{1}{2}(\delta(\omega + \omega_0) + \delta(\omega - \omega_0))$ ,  $\omega_0 \in \mathbb{R}$ .

5.4. **Odd Combination of Dirac-Deltas.**  $\mathfrak{F}^{-1}\{\hat{f}\}$  where  $\hat{f}(\omega) = \frac{1}{2i}(\delta(\omega + \omega_0) - \delta(\omega - \omega_0))$ ,  $\omega_0 \in \mathbb{R}$ .

5.5. **Horizontal Shifts.** Find  $\hat{f}(\omega)$  where  $f(x + c)$ ,  $c \in \mathbb{R}$ .

5.6. **Triangle Functions.** Find  $\mathfrak{F}\{f\}$  where  $f(x) = \begin{cases} 1 - |x|, & -1 < x < 1, \\ 0, & \text{otherwise} \end{cases}$

<sup>5</sup>A similar transformation can be found for the wave equation with inhomogeneous boundary conditions. The moral here is that time-dependent boundary conditions can be transformed into externally driven (AKA Forced or inhomogeneous) PDE with standard boundary conditions.

<sup>6</sup>Here the  $\delta$  is the so-called Dirac, or continuous, delta function. This isn't a function in the true sense of the term but instead what is called a generalized function. The details are unimportant and in this case we care only that this Dirac-delta *function* has the property  $\int_{-\infty}^{\infty} \delta(x - x_0)f(x)dx = f(x_0)$ . For more information on this matter consider [http://en.wikipedia.org/wiki/Dirac\\_delta\\_function](http://en.wikipedia.org/wiki/Dirac_delta_function). To drive home that this *function* is strange, let me spoil the punch-line. The sampling function  $f(x) = \text{sinc}(ax)$  can be used as a definition for the Delta *function* as  $a \rightarrow 0$ . So can a bell-curve probability distribution. Yikes!